

Disruption of gene expression in hybrids of the fire ants *Solenopsis invicta* and *Solenopsis richteri*

LINO OMETTO,*§ KENNETH G. ROSS,† DEWAYNE SHOEMAKER‡ and LAURENT KELLER*
*Department of Ecology and Evolution, University of Lausanne, Biophore, 1015 Lausanne, Switzerland, †Department of Entomology, University of Georgia, Athens, GA 30602-2603, USA, ‡Center for Medical, Agricultural, and Veterinary Entomology, United States Department of Agriculture – Agricultural Research Service, Gainesville, FL 32608, USA

Abstract

Transcriptome analysis is a powerful tool for unveiling the distribution and magnitude of genetic incompatibilities between hybridizing taxa. The nature of such incompatibilities is closely associated with the evolutionary histories of the parental species and may differ across tissues and between the sexes. In eusocial insects, the presence of castes that experience divergent selection regimes may result in additional distinct patterns of caste-specific hybrid incompatibilities. We analysed levels of expression of >14 000 genes in two life stages of each caste in the fire ants *Solenopsis invicta* and *Solenopsis richteri* and in their hybrids. We found strong contributions of both developmental stage and caste to gene expression patterns. In contrast, variability in expression was only weakly associated with taxonomic identity, with hybrid scores falling between those of the two parental species. Hybrid incompatibilities were surprisingly modest, with only 32 genes being mis-expressed, indicating low levels of disruption in gene regulation in hybrids; males and workers each mis-expressed at least seven times as many genes as queens. Interestingly, homologues of many of the mis-expressed genes have been implicated in behavioural variation in *Drosophila melanogaster*. General expression profiles of hybrids consistently were more similar to those of *S. richteri* than *S. invicta*, presumably because *S. richteri* trans-regulatory elements tend to be dominant and/or because there is an overall bias in the genetic composition of the hybrids towards *S. richteri*. Altogether, our results suggest that selection acting on each caste may contribute differently to interspecific divergence and speciation in this group of ants.

Keywords: ants, caste, gene expression, hybrids, *Solenopsis invicta*, *Solenopsis richteri*

Received 20 November 2011; revision received 27 January 2012; accepted 8 February 2012

Introduction

The process of speciation in eukaryotes typically involves divergence of two lineages from an ancestral lineage, resulting in their genomes becoming sufficiently incompatible that hybridization produces inviable or relatively less fit individuals. Such divergence can involve variable numbers of loci and may stem from neutral stochastic pressures or from extrinsic or intrinsic selective forces (Coyne & Orr 2004). For

instance, when species-specific loci have evolved to confer adaptation to divergent niches, hybrids may suffer fitness losses from having an intermediate phenotype poorly suited to either of the parental habitats (Schluter 2001; Taylor *et al.* 2012). Similarly, reduced hybrid fitness may derive from deleterious epistatic interactions between divergent parental alleles (i.e. Dobzhansky–Muller incompatibilities, Dobzhansky 1937; Muller 1942). These latter interactions typically involve co-adapted gene complexes and networks, implying that incompatibilities can involve both structural genes and regulatory regions. Consequently, accumulation of species-specific genetic variation can result in divergent regulatory networks that are disrupted in introgressed genomes, causing profound alterations in gene expres-

Correspondence: Lino Ometto, Fax: +39 0461 650956;

E-mail: lino.ometto@fmach.it

§Present address: Department of Biodiversity and Molecular Ecology, Research and Innovation Centre, Fondazione Edmund Mach, 38010 San Michele all'Adige (TN), Italy.

sion profiles compared to the parental species (e.g. Ranz *et al.* 2004; Landry *et al.* 2007; Ortiz-Barrientos *et al.* 2007; Renaut & Bernatchez 2011).

The genomic distribution and magnitude of such hybrid incompatibilities are influenced by the degree of genetic divergence at particular genomic elements, the effect sizes of diverging elements, dominance relationships among the parental alleles and patterns of epistasis (e.g. Orr 1993a; Turelli & Orr 1995). Thus, hybrid incompatibilities are tied to the evolutionary histories of the interacting genes in the parental species. Because the selective history of a gene is linked to its expression pattern and, ultimately, its phenotypic effect, the degree of incompatibility also may differ among environments, across tissues, and/or between the sexes. An additional layer of phenotypic differentiation often occurs in social hymenopteran insects as a result of the caste system, whereby females conditionally develop into fertile queens that are morphologically, physiologically and behaviourally distinct from sterile workers.

In highly eusocial Hymenoptera such as most ants and the honey bee, the striking behavioural and morphological differences between queens, workers and males are associated with important differences in historical and current selection regimes acting on each caste (Hunt *et al.* 2010, 2011; Ometto *et al.* 2011). Not surprisingly, previous studies have revealed that differences between social insect castes are associated with striking differences in gene expression during immature development (Evans & Wheeler 2000; Scharf *et al.* 2003; Pereboom *et al.* 2005; Hoffman & Goodisman 2007; Weil *et al.* 2009) as well as in the adult stage (Judice *et al.* 2006; Sumner *et al.* 2006; Gräff *et al.* 2007; Sen Sarma *et al.* 2007; Hunt & Goodisman 2010; Colgan *et al.* 2011; Ometto *et al.* 2011). Thus, interspecific hybridization may result in distinct caste-specific and sex-specific patterns and levels of hybrid incompatibilities in these insects. Accordingly, comparison of the nature of incompatibilities between the two female castes is expected to provide unique information on the relative importance of sexual (queen) versus somatic (worker) divergence in driving hybrid incompatibilities and reproductive isolation. Notably, incompatibilities arising solely in workers have only an indirect effect on hybrid fitness, because members of this caste generally do not reproduce, yet they profoundly affect colony survival and productivity.

Hybridization in the wild has been well documented in several species of ants (e.g. Pearson 1983; Douwes & Stille 1991; Shoemaker *et al.* 1996; Seifert 1999; Helms Cahan *et al.* 2002; Feldhaar *et al.* 2003; Helms Cahan & Vinson 2003; Umphrey 2006; Kulmuni *et al.* 2010; reviewed in Feldhaar *et al.* 2008). However, excluding unusual reproductive systems such as those in which

historical hybridization has led to the stable coexistence of distinct interbreeding lineages (Helms Cahan & Keller 2003; Helms Cahan *et al.* 2004; Schwander *et al.* 2006), very few studies have investigated any of the phenotypic or fitness consequences of hybridization in ants (Jessen & Klinkicht 1990; Ross & Robertson 1990; Pusch *et al.* 2006). Thus, the genetic basis of hybrid incompatibilities manifested in each caste, and the contributions of these incompatibilities to reproductive isolation between hybridizing species, remains largely unknown for ants as well as other social insects.

A valuable study system for investigating patterns of hybrid incompatibility in social Hymenoptera is represented by the closely related fire ant species *Solenopsis invicta* and *Solenopsis richteri*. These two species are reproductively isolated in their native South American ranges (Ross & Shoemaker 2005) but exhibit extensive introgression and evidently produce relatively fit hybrids in their invasive North American ranges (Ross & Robertson 1990; Shoemaker *et al.* 1996). The absence of premating reproductive isolation between the USA populations may stem in part from the fact that they originated from allopatric South American populations that lack the behavioural isolating mechanisms produced by reinforcement in areas of sympatry in their native ranges. Alternatively, specific ecological or environmental features constituting barriers to hybridization in the native ranges may be lacking in the introduced range, thus compromising reproductive isolation caused by extrinsic factors. Finally, the genetic bottlenecks that each species experienced upon introduction to the USA may have compromised genetically based mate recognition systems that enforce reproductive isolation in the native ranges. Regardless of the nature of the premating barriers that have been compromised, the fact that postmating barriers between the species are incompletely developed provides the opportunity for in-depth study of the genomics of hybrid incompatibility in a wild setting.

Previous genetic, biochemical and morphological studies of the large zone of hybridization between *S. invicta* and *S. richteri* in the USA revealed that it features a gradual changeover in allele frequencies and character sizes from those typical of *S. invicta* in the south to those typical of *S. richteri* in the north (Ross *et al.* 1987; Ross & Robertson 1990; Shoemaker *et al.* 1996). Highly admixed hybrid genotypes predominate in the centre of the zone, consistent with the occurrence of hybridization over many generations. Moreover, genotypes in these highly admixed populations typically occur at frequencies expected under Hardy–Weinberg equilibrium, and overall significant pairwise linkage disequilibria are effectively absent. Although large-scale clinal changes in genetic and phenotypic characters are the norm

along study transects spanning the zone, pockets of pure parental forms and hybrids of varying genetic constitution are interspersed throughout it; this mosaic small-scale distribution may reflect chance historical colonization of suitable habitat patches or preferential nesting success of hybrids of varying genomic composition among ecologically distinct microhabitats (Shoemaker *et al.* 1996; the parental species occupy different habitats in their native ranges – Trager 1991). The ubiquity of advanced-generation hybrid genotypes in the USA hybrid zone, combined with the general lack of single-locus and linkage disequilibria, suggests not only that F1 hybrids are viable and fertile, but also that any general breakdown in hybrid fitness may be modest.

The possibility that hybrid incompatibility may cause disrupted development in hybrid *S. invicta/S. richteri* was earlier studied by comparing the developmental stability of several queen morphological characters between hybrids and the parental species (Ross & Robertson 1990). Hybrid ants displayed significantly reduced developmental stability, but the effect was weak. Thus, genomic incompatibility between the parental species may create only modest viability selection against hybrid genotypes that is not sufficient to preclude extensive hybridization. Moreover, the finding that some diagnostic genetic markers introgress more freely across the hybrid zone than species-typical morphological traits (Shoemaker *et al.* 1996) suggests that genes underlying the latter recombine less readily into a heterospecific genomic background, consistent with mild intrinsic fitness penalties to at least some recombinant genotypes.

A previous study of genome-wide expression in *S. invicta* and *S. richteri* revealed low divergence in the expression patterns of the two species (Ometto *et al.* 2011). However, this differentiation varies greatly between castes and developmental stages, with the largest differences in gene expression occurring between adult workers of the two species. The aim of this study was to determine levels of gene expression in hybrids to learn whether the various castes, sexes and life stages also differ in the degree of compatibility between the *S. invicta* and *S. richteri* regulatory machineries. In particular, we were interested in investigating whether differences in life history between castes result in greater numbers of genes being mis-expressed in workers than in males and queens. Because fire ants workers do not reproduce, and thus have only an indirect fitness component, gene expression may be under more relaxed selection in this caste than in queens and males (Linksvayer & Wade 2009; Ometto *et al.* 2011). If we assume that regulation of gene expression has a genetic basis and is under stronger purifying selection in reproductively competent individuals, genes preferentially

expressed in workers should be regulated by elements exhibiting a greater between-species divergence than elements imposing similar gene expression levels across castes. Moreover, there should also be relatively greater interspecific divergence in the patterns of selection acting on workers; this is because adult members of this caste most directly experience the distinct environments characterizing the different ranges occupied by the two species (namely, tropical and subtropical habitats for *S. invicta* and somewhat more temperate habitats for *S. richteri* – Trager 1991), whereas queens and males are almost exclusively confined to the buffered, within-nest environment. As a result, we predict that there should be more genes mis-expressed in workers than in queens and males. Similarly, pupae are entirely confined within the nest, where they are cared for by the workers, and therefore, they presumably also experience more similar environments across the two species than do adults, leading to the prediction that pupae should have fewer genes mis-expressed than adults.

Materials and methods

We analysed gene expression levels of workers, queens and males in the pupal and adult stages of *S. invicta* X *S. richteri* hybrids as well as the two parental species. As typically is the case in ants, it is impossible to conduct controlled matings with fire ants under laboratory conditions. Therefore, colonies of hybrids were collected directly from the field in Grenada and Alcorn Counties, Mississippi, USA (a central location in the hybrid zone) in May of 2005 and 2007. Upon return to the laboratory, hybrid as well as pure-species colonies were maintained in a single rearing room under identical standard conditions, including the same artificial nests, diets, and temperature and humidity regimes (Jouvenaz *et al.* 1977).

Taxonomic status of every study colony was confirmed by genotyping 8–10 individuals from each at five informative allozyme loci that feature strong or fixed allelic differences between the species (see Shoemaker *et al.* (1996) for details of the allozyme methods).

We conducted analyses of gene expression patterns in hybrids as previously described for pure *S. invicta* and *S. richteri* (Ometto *et al.* 2011). Briefly, we randomly sampled, and pooled, 1–6 nestmate individuals (mean = 4.6, median = 5.0) for each replicate of the eighteen specific categories, or nodes, of interest (two developmental stages X 3 castes X 3 taxa (two species + one hybrid); Table S1, Supporting information). Specimens of both developmental stages were standardized by age (pupae – first appearance of pink eyes; adults – <12 h posteclosion), and specimens of the worker caste were further standardized by including only majors so as to minimize possible allometric

differences associated with intracaste polyphenism, which could affect gene expression levels. The cDNA derived from each experimental sample was hybridized against a common reference RNA on our custom-made spotted cDNA microarray after being randomly labelled with either Cy3 or Cy5 (Wang *et al.* 2007; Wurm *et al.* 2009). Because we used each colony only once for a given caste and developmental stage, each sample was equivalent to an independent biological replicate. Microarray intensity analyses were performed using the software package *limma* in R (R Development Core Team 2009). The power of the between-array normalization was enhanced by adding data from arrays hybridized with cDNA originating from larvae, which underwent the same experimental protocols as the samples used in the present study and in Ometto *et al.* (2011). Raw data for all 140 hybridizations (i.e. pure species and hybrid data for workers, queens and males at the larval, pupal and adult stages) are available in the Gene Expression Omnibus database under accession number GSE35217.

The Bayesian approach implemented in the program Bayesian Analysis of Gene Expression Levels (BAGEL; Meiklejohn & Townsend 2006) was used to estimate the relative expression level of each clone for each node (stage/caste/taxon) and the significance of the differential expression of such clones among nodes (data deposited in the Dryad repository: doi:10.5061/dryad.m0r5qv24). Significance was estimated after correcting for multiple testing using the false discovery rate approach as described in Ometto *et al.* (2011). If at least one of the clones belonging to the same contig was differentially expressed, the contig was considered differentially expressed as well. For consistency with our previous study (Ometto *et al.* 2011), we report only results based on contigs, and we use contig synonymously with gene throughout the remainder of this text. Following Ometto *et al.* (2011), a given gene was categorized as caste-biased in its expression when significantly over- or under-expressed in one caste compared to both of the other castes for a given developmental stage.

Some regulatory mechanisms/factors that have diverged between *S. invicta* and *S. richteri* may not necessarily translate into different gene expression levels or different phenotypes between the two species (True & Haag 2001). However, such divergence could be apparent in hybrids that contain a mix of different and possibly incompatible regulatory components (Riddle & Birchler 2003), resulting in consistently over- or under-expressed transcripts relative to the pure parental species (e.g. Michalak & Noor 2003; Ranz *et al.* 2004; Haerty & Singh 2006). We therefore quantified the degree of hybrid incompatibilities in gene regulation by compar-

ing the level of expression of each gene between hybrids and each of the two parental species. Genes were defined as mis-expressed when either significantly over- or under-expressed in hybrids compared to both parental species. These comparisons were performed separately for each developmental stage and caste.

The relative expression of each contig (measured as the average of the values calculated by BAGEL for the associated clones) was used to estimate the relative importance of developmental stage, caste and taxonomic identity on the observed gene expression levels across all genes. First, we estimated pairwise gene expression distances between each of the eighteen nodes and constructed the corresponding distance tree as described in Ometto *et al.* (2011). In a second approach, we performed principal components analyses on the scaled data matrix using the singular value matrix decomposition approach of the *prcomp* package implemented in R (R Development Core Team 2009).

The normalized gene expression data were used to estimate the variability in gene expression, V_x , within *S. invicta*, *S. richteri*, and hybrids, and to evaluate H_{ir} , the degree of similarity in gene expression between hybrids and the two parental species. Specifically, V_x was estimated as the ratio between the mean and the standard deviation of the \log_2 ratios of the normalized dye intensities calculated across all samples for the species, caste and developmental stage of interest (see Ometto *et al.* (2011) for details of the methods). The similarity index H_{ir} was estimated for each clone and hybrid sample as, $(\left|\frac{h-i}{i}\right| - \left|\frac{h-r}{r}\right|)$, where h is the normalized gene expression for the sample of interest, and i and r are the across-samples mean expression values for *S. invicta* and *S. richteri*, respectively.

We used gene ontology (GO) terms to test whether genes differentially expressed between hybrids and the parental species were over-represented (enriched) for some categories. Because no GO terms are available for *S. invicta*, we used the Blast2GO web tool (Conesa *et al.* 2005) to functionally annotate the *S. invicta* clone sequences available from Fourmidable, <http://fourmidable.unil.ch> (Wang *et al.* 2007; Wurm *et al.* 2009). Searching within the National Center for Biotechnology Information (NCBI) database, the tool could assign a putative orthologue to 10 527 clones, 6197 of which had associated GO terms. In a second approach, mapping information (Hunt *et al.* 2011) allowed us to assign 4767 of the 22 856 clones spotted onto the microarray to 2066 *S. invicta* genes (official gene set version 2.2.0, available from Fourmidable, <http://fourmidable.unil.ch>; Wurm *et al.* 2009, 2011). We then searched for putative orthologues in the well-annotated *Drosophila melanogaster* genome (official gene set release 5.9, available at <http://flybase.org> – Tweedie *et al.* 2009).

Orthology was determined by BLASTp searches (Camacho *et al.* 2009), and custom Perl scripts were used to identify a total of 1403 reciprocal best hits, which were retained as pairs of putative orthologues. The annotations of the *D. melanogaster* orthologues were subsequently used to search for potential enrichment by GO terms using the DAVID bioinformatics tool (Huang *et al.* 2009). Functional prediction based on orthology may be compromised when conducted with distantly related species (e.g. fruit flies and fire ants) at the single gene level (e.g. Nehrt *et al.* 2011), but this potential problem should only minimally affect global conclusions.

Results and Discussion

Principal components analysis revealed a strong contribution of developmental stage to gene expression, with the first component, on which this variable loads heavily, explaining 42% of the total gene expression variability (Fig. S1, Supporting information). The second, third and fourth components, associated with caste and sex, together accounted for 39% of the expression variability. The fifth component, explaining 5% of the gene expression variability, was not distinctly associated with any of the variables. Only the sixth component, which accounts for just 4.2% of the expression variability, was associated with taxonomic identity, with hybrid scores falling between the two parental species' scores in both life stages of all castes (Fig. S1, Supporting information). The fact that taxonomic group explains so little of the variability in the expression data implies strong conservation in gene expression patterns between the two parental species. This conclusion also is supported by the close clustering of hybrids with the two parental species within each stage and caste in the gene expression-based distance tree (Fig. 1).

Gene expression differentiation between hybrids and the parental species

Explicit comparisons of gene expression levels between hybrids and the two parental species support the view that the conserved expression patterns translate into limited incompatibilities in gene regulation between *S. invicta* and *S. richteri*. In male pupae, there were only 147 genes (1.0% of the 14,467 genes analysed) that were differentially expressed between hybrids and either of the two parental species (Fig. 2). In female pupae, the numbers were similarly low, with 239 (1.7%) genes differentially expressed in hybrid queens and 319 (2.2%) in hybrid workers. In adults, there were slightly more genes differentially expressed between hybrids and the parental species, with 310 (2.1%) such genes in males, 247 (1.7%) in queens and 662 (4.6%) in workers (Fig. 2).




Fig. 1 Gene expression-based distance tree. Three-letter labels identify taxonomic group (i = *Solenopsis invicta*; r = *Solenopsis richteri*; h = hybrids), caste (W = workers; Q = queens; M = males) and developmental stage (p = pupae; a = adults).

When considering all three castes in aggregate for the pupal stage, 4.0% of genes were differentially expressed between hybrids and *S. invicta* and 1.1% between hybrids and *S. richteri*. A similar comparison previously revealed that 5.8% of genes were differentially expressed between pupae of the two parental species (Ometto *et al.* 2011). In adults, the patterns were similar, with 5.4% of genes differentially expressed in the hybrid-*invicta* comparison, 3.5% in the hybrid-*richteri* comparison and 10.7% in the *invicta-richteri* comparison. Importantly, our finding of a lower percentage of genes being differentially expressed between hybrids and either parental species than between the two species is conservative, because the false discovery rate for the hybrid *vs.* parental species analyses is at least twice as high as the false discovery rate for the interspecific analyses for both life stages (data not shown; Ometto *et al.* 2011).

This finding of much lower proportions of genes being differentially expressed between hybrids and either parental species than between the species (about 1/2) suggests relatively few incompatibilities and thus quite recent divergence between *S. invicta* and *S. richteri*. For comparison, approximately equal numbers of genes are differentially expressed between F1 *Drosophila simulans* X *Drosophila sechellia* hybrids and the parental species as between the two species (Haerty & Singh 2006; Moehring *et al.* 2007; Wurmser *et al.* 2011), which evidently separated some 250 000 years ago (Mcdermott & Kliman 2008). Moreover, about twice as many genes are differentially expressed between F1 *Drosophila melanogaster* X *D. simulans* hybrids and the parental species as between the two species (Ranz *et al.* 2004), which are




Fig. 2 Venn diagrams depicting numbers of genes significantly overexpressed (>) or under-expressed (<) in pupal and adult hybrids (hyb) relative to *Solenopsis invicta* (inv) and *Solenopsis richteri* (ric). Blue-shaded overlapping areas identify genes consistently over- or under-expressed in hybrids relative to both parental species (i.e. mis-expressed genes). Grey-shaded overlapping areas identify genes that are significantly over-expressed in hybrids compared to one parental species while significantly under-expressed compared to the other. Sizes of circles are proportional to the numbers of genes.

estimated to have diverged about 2–3 million years ago (Hey & Kliman 1993). While direct extrapolation of times of divergence across these insect orders is not appropriate, the fire ant regulatory element divergence clearly is comparatively low. This is in line with other evidence that the clade to which our two study species belong (the *Solenopsis saevissima* species group) is a youthful group in an active phase of evolutionary radiation (Ross & Trager 1990; Shoemaker *et al.* 2005).

One factor complicating explanations of the relatively low divergence in gene expression levels between the fire ant hybrids and the two parental species is that the sampled hybrids were not F1 hybrids, but rather a mix of various advanced-generation hybrids and backcross-

es. Thus, individuals (colonies) with high levels of hybrid incompatibilities may have been selectively purged in the wild and thus unavailable for sampling. While this effect may contribute to an underestimate of the extent of transcriptome divergence between *S. invicta* and *S. richteri*, such estimates are most relevant to the ecological context of natural hybridization. Clearly, the ecologically effective level of genetic incompatibilities between the two parental genomes is sufficiently low that (i) hybrids are widespread and abundant in the introduced (USA) range; (ii) introgression has proceeded through multiple generations to yield highly admixed recombinant genotypes that predominate in the hybrid zone centre; and (iii) developmental stability has been little disrupted in extant hybrid queens (Ross & Robertson 1990; Shoemaker *et al.* 1996).

Mis-expressed genes in hybrids

Consistent with the finding of low divergence in gene expression profiles between hybrids and the parental species, we also found low levels of disruption in gene regulation in hybrids. There were only 16 mis-expressed genes (i.e. genes significantly over- or under-expressed in hybrids compared to both parental species) in pupae and 18 in adults (Fig. 2), only one of which was mis-expressed in both stages. When analysed individually, however, the three castes differed greatly in the number of genes mis-expressed, with the male and worker castes each mis-expressing at least seven times more genes than the queen caste in both life stages (goodness-of-fit test comparing the observed data to the expectation of equal numbers of mis-expressed genes across castes; $P < 0.05$ for all such comparisons; $P > 0.8$ for the comparisons between males and workers for both stages). The haploid genomic state of males may contribute to the large number of mis-expressed genes in this caste, because haploidy should result in admixed males expressing all recessive incompatibilities after the F1 female generation (i.e. after a first round of meiotic recombination between the paternal and maternal homologues; Haldane 1922; Orr 1993b; Turelli & Orr 1995; Schilthuizen *et al.* 2011; Koevoets *et al.* 2012). On the other hand, the significant difference in the number of mis-expressed genes between the two female castes can only be explained by higher levels of regulatory incompatibilities in hybrid workers than in hybrid queens. This finding is in line with the prediction that, because workers are sterile and experience more divergent environments than queens and males, they should exhibit higher levels of regulatory incompatibilities (see Introduction). The observed lower number of differentially expressed (as opposed to mis-expressed) genes in hybrid pupae than in hybrid adults, particularly notable

in males and workers (Fig. 2), agrees with the greater conservation of gene expression profiles observed in pupae of the pure species compared to adults (Ometto *et al.* 2011); these patterns are consistent with pupae experiencing a more buffered environment than adults as well as with a generally greater conservation of regulatory networks involved in early than in late development during cladogenesis (Artieri & Singh 2010; Domazet-Lošo & Tautz 2010).

One potentially confounding factor in interpreting patterns of mis-expression based on whole-body mRNA extracts, such as used in our study, is the possible atrophy of various tissue/organ systems owing to hybrid breakdown, one result of which may be differences in mRNA abundance attributable solely to gross differences in tissue abundance between hybrids and pure-species individuals (e.g. Ranz *et al.* 2004; Blumenstiel & Hartl 2005). While we cannot entirely rule out a role for such effects, several lines of evidence suggest they may not be important. First, these effects are not expected to produce the commonly observed pattern of hybrid fire ant expression exceeding parental expression, barring the occurrence of hypertrophy rather than atrophy of hybrid tissues/organs. Second, fire ant workers effectively lack a reproductive system (Tschinkel 2006), so that the atrophy of reproductive tissues in hybrids commonly underlying artifactual expression differences between them and parentals (Hollocher *et al.* 2000; Ranz *et al.* 2004; Graveley *et al.* 2011) is not possible in this caste. Finally, our hybrid study samples did not generally exhibit greater variation in expression levels across replicate colonies than did the parental species, as would be predicted given that the hybrids represent various levels of backcrossing and assuming that this would yield corresponding variation in tissue dystrophy. Among the six developmental stage/caste combinations, the degree of gene expression variability, V_X , in hybrids significantly surpassed that in both parental species in only half of the cases (pupal workers and males and adult queens; Wilcoxon test, all $P < 0.001$), suggesting minimal general effects of any hybrid tissue atrophy/hypertrophy on expression differences.

Functional characterization of mis-expressed genes

Functional annotation of the 33 mis-expressed genes was conducted by searching for their putative orthologues and related GO terms in available annotated genomes. In a first approach, we probed the sequences of the clones associated with the mis-expressed genes against the National Center for Biotechnology Information database using the Blast2GO tool (Conesa *et al.* 2005). We could assign putative orthologues to 14 of the 33 mis-expressed genes, most of which matched genes

from ants (Table 1) (Bonasio *et al.* 2010; Nygaard *et al.* 2011; Wurm *et al.* 2011). It was possible to associate GO terms with only five of these genes, and no significant over-representation of any functional class was apparent in these annotation data. The limited success of our functional annotation efforts may stem from the fact that clone sequences represent only partial genes, which reduces the efficiency of the orthologue searches.

In a second approach specifically aimed at overcoming this limitation and targeting well-annotated genomes, we were able to identify seven of the 33 mis-expressed genes by reference to the *S. invicta* genome, and five of these were inferred to have an identifiable *D. melanogaster* orthologue (Table 1). Two of the *S. invicta* genes matched the same *D. melanogaster* orthologue, suggesting a possible duplication event in an ancestral fire ant lineage and reducing the number of putative orthologue pairs to four. Two of these four genes are annotated in *D. melanogaster* as being involved in nitrogen compound biosynthetic processes, a significant over-representation of this gene class according to the gene ontology analysis of the *S. invicta*-*D. melanogaster* orthologue pairs (Fisher's exact test; $P = 0.03$).

Interestingly, four of the genes for which we could infer putative functions have been implicated as being involved in behavioural variation in *D. melanogaster* (Table 1). For instance, gene CG3011 is related to the response to ethanol exposure (Kong *et al.* 2010), while gene *ade5* is involved in between-male aggression (Edwards *et al.* 2009). The *S. invicta* orthologues of both genes were under-expressed in hybrids compared to both parental species in adult workers. The third gene codes for the cuticular protein 72Ec, while the fourth gene is a putative homologue of the odorant/hormone-binding gene CG14661 in *D. melanogaster*. Both of these genes show circadian oscillations in *D. melanogaster* (Ceriani *et al.* 2002; Wijnen *et al.* 2006), and in *Solenopsis*, they were overexpressed in hybrids compared to both parental species (*Cpr72Ec* in male pupae and CG14661 in worker pupae). Given the paramount importance of behavioural interactions in the social lives of ants, both between adults and between pupae and nursing adult workers, mis-expression of genes influencing the performance of, or responses to, behaviours can be expected to have profound fitness consequences in hybrid fire ants not necessarily evident at the physiological or morphological levels (cf. Ross & Robertson 1990).

Asymmetries in hybrid gene expression profiles

The expression profiles of hybrids in both developmental stages were significantly more similar to those of

Table 1 Putative orthologues of the genes mis-expressed in hybrid fire ants

Name*	Mis-type [†]	Blast2GO [‡]		Gene	Gene ontology (GO) terms/function	Gene	GO terms/function
		<i>Solenopsis invicta</i> gene	<i>Drosophila melanogaster</i> [§]				
SjJWA08BAL	+aM	NA	Hypothetical protein G5I_14350 (<i>Acromyrmex echinatior</i>)	NA	Binding	NA	NA
SjJWB02BCV	-aW	NA	NA	NA	NA	NA	NA
SjJWB10AEA	-aW	NA	NA	NA	NA	NA	NA
SjJWC03ACC	+aM	NA	Alpha-tocopherol transfer (<i>S. invicta</i>)	NA	Transporter activity	NA	NA
SjJWC04BCM	+pW	NA	NA	NA	NA	NA	NA
SjJWC04BDK	-pM	NA	NA	NA	NA	NA	NA
SjJWC07ADV	-pM	SI2.2.0_06746	NA	CG14661	NA	CG14661	Odorant/hormone-binding protein involved in circadian rhythm
SjJWC10CAM	+aM	SI2.2.0_05389	Hypothetical protein SINV_05389 (<i>S. invicta</i>)	NA	NA	NA	NA
SjJWD04BCT	-pM	NA	NA	NA	NA	NA	NA
SjJWD07BEA	-pM	SI2.2.0_06746	Circadian clock-controlled protein (<i>S. invicta</i>)	CG14661	NA	CG14661	Odorant/hormone-binding protein involved in circadian rhythm
SjJWD10ACV	+aM	NA	Transmembrane protein 39A-A-like (<i>Acromyrmex echinatior</i>)	NA	NA	NA	NA
SjJWD10BAK	+pW	NA	NA	NA	NA	NA	NA
SjJWE06BAD	+aW	NA	NA	NA	NA	NA	NA
SjJWE10AAV	-aW	SI2.2.0_07244	Multifunctional protein ade2 (<i>S. invicta</i>)	Multifunctional protein <i>ade5</i>	Nitrogen compound biosynthetic process	Multifunctional protein <i>ade5</i>	Nitrogen compound biosynthetic process; involved in intermale aggression
SjJWE12BAI	+pM	NA	NA	N	N	N	N
SjJWE12BDW	+aW	NA	NA	N	N	N	N
SjJWF01ABR	-aQ	NA	NA	N	N	N	N
SjJWF01BBB	-aW	SI2.2.0_09632	Serine hydroxymethyltransferase (<i>S. invicta</i>)	CG3011	L-serine metabolic process; methylation; nitrogen compound biosynthetic process.	CG3011	Serine hydroxymethyltransferase; nitrogen compound biosynthetic process; related to the response to ethanol exposure
SjJWF08BBH	-aM	NA	NA	NA	NA	NA	NA

Table 1 Continued

Name*	Mis-type [†]	Blast2GO [‡]		<i>Drosophila melanogaster</i> [§]		
		<i>Solenopsis invicta</i> gene	Gene	Gene ontology (GO) terms/function	Gene	GO terms/function
SIJWF09BDN	-aW	NA	NA	NA	NA	NA
SIJWG01BBA	-aW	NA	NA	NA	NA	NA
SIJWG04AAG	-aW	NA	NA	NA	NA	NA
SIJWG06CAG	pM	SI2.2.0_01800	PR domain zinc finger protein 13 (<i>S. invicta</i>)	Zinc ion binding; nucleic acid binding		
SIJWG09BCP	-pW	SI2.2.0_13401	Cuticle protein 6 (<i>S. invicta</i>)	NA	<i>Cpr72Ec</i> (Cuticular protein 72Ec7)	Shows circadian oscillations
SIJWH02BCR	-aM	NA	Hypothetical protein SINV_00721 (<i>S. invicta</i>)	NA	NA	NA
SIJWH03AAT	-pQ	NA	Hypothetical protein	NA	NA	NA
SIJWH04AAK	-pW	NA	Hypothetical protein SINV_16046 (<i>S. invicta</i>)	NA	NA	NA
SIJWH04ACW	-pW	SI2.2.0_04262	Hypothetical protein EAI_10442 (<i>Harpegnathos saltator</i>)	NA	NA	NA
SIJWH05BCO	-p/aM	NA	Chymotrypsin-2 (<i>S. invicta</i>)	NA	NA	NA
SIJWH06CAH	-aM	NA	NA	NA	NA	NA
SIJWH07ACG	-pM	NA	WD-repeat-containing protein 16 (<i>S. invicta</i>)	NA	NA	NA
SIJWH09ABR	-pW	NA	NA	NA	NA	NA
SIJWH09ADK	-pW	NA	NA	NA	NA	NA

NA, not available.

*Name of the clone spotted onto the microarray (Wurm *et al.* 2009).

[†]Type of mis-expression pattern of the clone. The three-letter code indicates whether a clone was overexpressed (+) or under-expressed (-) in hybrids compared to both parental species in pupal (p) or adult (a) workers (W), queens (Q), or males (M).

[‡]Putative orthologues of the clones and associated GO terms.

[§]Putative orthologues of the *S. invicta* genes identified by a reciprocal best hit approach in *D. melanogaster* and associated GO terms.

S. richteri than *S. invicta* (goodness-of-fit test, both $P < 10^{-13}$). In pupae, the degree of hybrid expression bias towards *S. richteri* was strong in all three castes (Fig. 2). In queens of this stage, more than twice as many genes were differentially expressed between hybrids and *S. invicta* than between hybrids and *S. richteri* (177 vs. 70; goodness-of-fit test; $P < 10^{-11}$), while in males and workers, there were four times more genes differentially expressed between hybrids and *S. invicta* than between hybrids and *S. richteri* (127 vs. 28 for males and 276 vs. 56 for workers; both $P < 10^{-14}$). In the adult stage, however, there were important differences among the three castes (Fig. 2). While hybrid expression biases towards *S. richteri* were similar to those in pupae for both adult queens (214 vs. 41; $P < 10^{-15}$) and adult males (261 vs. 617; $P < 10^{-15}$), the expression bias in hybrid adult workers was towards *S. invicta*, with more genes differentially expressed between hybrids and *S. richteri* than between hybrids and *S. invicta* (398 vs. 300; $P < 10^{-15}$). The distinctive gene expression profile of hybrid adult workers, which departed from the overall pattern of bias towards *S. richteri*, was further examined by two additional analyses.

In the first analysis, we compared the ratio of expression for hybrids and *S. richteri* to the ratio for hybrids and *S. invicta* (Fig. 3). The expression levels in hybrid pupae of each of the three castes were closer to those of *S. richteri* than to those of *S. invicta* (paired Wilcoxon test; $P < 10^{-11}$ for all three castes). The same pattern was found for adult queens ($P < 10^{-10}$) but was reversed for adult males ($P < 10^{-10}$). No significant pattern emerged for adult workers ($P = 0.383$).

In the second analysis, we examined the degree of conservation of caste-biased gene expression between hybrids and each parental species by comparing the numbers of genes over- and under-expressed in a par-

ticular caste in both hybrids and *S. invicta* to the numbers of genes over- and under-expressed in that same caste in both hybrids and *S. richteri*. In pupae, genes that were caste-biased in hybrids were significantly more likely to be caste-biased as well in *S. richteri* than in *S. invicta* (Fisher's exact tests, for overexpressed genes $P < 0.0001$ for workers, queens and males; for under-expressed genes $P < 0.05$ for workers and queens and $P > 0.17$ for males; Figs S2a and S3a, Supporting information respectively). In adults, such differences, while apparent for each caste, were statistically significant only for overexpressed genes in males ($P < 10^{-9}$; $P > 0.183$ for overexpressed genes in workers and queens and for under-expressed genes in all three castes; Figs S2b and S3b, Supporting information). Overall, these results can be taken to indicate different interactions of the parental regulatory machineries both among castes and between developmental stages.

Two hypotheses may explain the finding that hybrids generally display gene expression patterns closer to *S. richteri* than to *S. invicta*. The first is directly linked to the relative contribution of *S. richteri* to the hybrid genome pool. The allozyme data revealed an overall bias in the genetic composition of the hybrid colonies towards *S. richteri*; assuming independence of the five loci analysed, the average hybrid index (Buerkle 2005) was 0.69 (range, 0.58–0.94 for the seven colonies), where a value of 0 corresponds to pure *S. invicta* and a value of 1 to pure *S. richteri*. Thus, it is probable that the hybrids analysed in this study had a larger fraction of their genomes originating from *S. richteri* than from *S. invicta*. In females, this would mean that an increased fraction of loci were homozygous for *S. richteri cis* or *trans* elements regulating the expression of associated genes compared to a population with a hybrid index of 0.5, and hence, the fraction of genes with expression patterns most similar to those of




Fig. 3 Ratios between the relative gene expression levels in hybrids and either *Solenopsis invicta* (inv) or *Solenopsis richteri* (ric). Values closer to one indicate greater similarity of gene expression levels between hybrids and a parental species. Error bars denote the standard error of the mean. Paired Wilcoxon test, **** $P < 0.0001$, NS = not significant.

S. richteri would exceed the fraction similar to *S. invicta*. To further test this hypothesis, we conducted correlation analyses between the colony-specific hybrid index and H_{irr} , an index of similarity in expression between hybrids and each parental species. Correlations were not significant ($P > 0.13$ for pupae and adults in all three castes). This indicates either that the genomic composition has only a marginal effect on the gene expression patterns of hybrids, or perhaps more likely, that the five allozyme loci in aggregate do not provide sufficient information to accurately predict the genome pool composition of individual colonies (and thus cannot be used to reliably test whether the bias in gene expression patterns of hybrids towards *S. richteri* stems from a higher contribution of *S. richteri* to the hybrid genome pool).

The second hypothesis is that *S. richteri* trans-regulatory elements are on average dominant to those of *S. invicta*, thus imposing an expression profile closer to that of *S. richteri* even with relatively even mixes of the two genomes. In support of this, the gene expression-based distance tree revealed similar overall divergence in gene expression patterns of hybrids from those of each parental species (Fig. 1), in contrast to the greater similarity of hybrids to *S. richteri* than *S. invicta* at the relatively small fraction of genes with significantly divergent expression in hybrids. Asymmetry in gene expression differences towards one of the parental species has also been observed in hybrids between sympatric anadromous and resident populations of brook charr (Mavarez *et al.* 2009) and in hybrids between lake whitefish species pairs (Renaut *et al.* 2009), pointing to the common existence of complex interactions between parental species' regulatory machineries when divergent genomes are admixed (Gibson & Weir 2005; Rockman & Kruglyak 2006).

Under both hypotheses, the asymmetry in expression profiles might be expected to hold across all three castes in both developmental stages, yet our results show that adult workers consistently differ in this regard from the other classes of individuals. This incongruity may stem from either of two nonmutually exclusive causes. First, adult worker gene expression may be under the control of regulatory elements that generally are more sensitive to genetic perturbations, such that expression levels in these individuals often differ from those of the parental species irrespective of the species origin of the regulatory elements. Alternatively, purifying selection may operate less efficiently in workers than in queens and males, resulting in delayed purging of incompatibilities in this caste as introgression proceeds (Linksvayer & Wade 2009). This explanation is appealing because workers are obligately sterile in these ants and so possess only indirect fitness components manifested by the

effects of their activities on the survival and reproduction of related reproductive nestmates.

Conclusion

This study reveals evidence of surprisingly modest hybrid incompatibilities in gene expression between two invasive fire ant species, *S. invicta* and *S. richteri*. This low level of incompatibility probably accounts in part for the widespread and persistent hybridization between the two species in the USA, where premating barriers have been compromised following their introductions in the early part of the last century. This study further highlights significant heterogeneity in the degree of hybrid incompatibilities across life stages and castes, including elevated numbers of genes mis-expressed in workers compared to males and queens and discordance among classes of individuals in the similarity of their gene expression profiles to those of each parental species. Altogether, these results suggest that selection pressures acting specifically on each caste may contribute differently to interspecific divergence and the processes of speciation in these ants as well as in other social insects.

Acknowledgements

We thank Christine La Mendola and the Lausanne Genomic Technologies Facility for technical support, the Keller laboratory for helpful discussions and two reviewers for helpful comments on the manuscript. This work was supported by grants from the Roche Research Foundation (to L.O.) and Swiss NSF (L.K.) as well as an ERC advanced grant (L.K.).

References

- Artieri CG, Singh RS (2010) Molecular evidence for increased regulatory conservation during metamorphosis, and against deleterious cascading effects of hybrid breakdown in *Drosophila*. *BMC Biology*, **8**, 26.
- Blumenstiel JP, Hartl DL (2005) Evidence for maternally transmitted small interfering RNA in the repression of transposition in *Drosophila virilis*. *Proceedings of the National Academy of Sciences of the United States of America*, **102**, 15965–15970.
- Bonasio R, Zhang G, Ye C *et al.* (2010) Genomic comparison of the ants *Camponotus floridanus* and *Harpegnathos saltator*. *Science*, **329**, 1068–1071.
- Buerkle C (2005) Maximum-likelihood estimation of a hybrid index based on molecular markers. *Molecular Ecology Notes*, **5**, 684–687.
- Camacho C, Coulouris G, Avagyan V *et al.* (2009) BLAST+: architecture and applications. *BMC Bioinformatics*, **10**, 421.
- Ceriani MF, Hogenesch JB, Yanovsky M *et al.* (2002) Genome-wide expression analysis in *Drosophila* reveals genes controlling circadian behavior. *The Journal of Neuroscience*, **22**, 9305–9319.

- Colgan TJ, Carolan JC, Bridgett SJ *et al.* (2011) Polyphenism in social insects: insights from a transcriptome-wide analysis of gene expression in the life stages of the key pollinator, *Bombus terrestris*. *BMC Genomics*, **12**, 623.
- Conesa A, Götz S, García-Gómez JM *et al.* (2005) Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics*, **21**, 3674–3676.
- Coyne JA, Orr HA (2004) *Speciation*. Sinauer Associates Inc, Sunderland, MA.
- Dobzhansky T (1937) *Genetics and the Origin of Species*. Columbia University Press, New York.
- Domazet-Lošo T, Tautz D (2010) A phylogenetically based transcriptome age index mirrors ontogenetic divergence patterns. *Nature*, **468**, 815–818.
- Douwes P, Stille B (1991) Hybridization and variation in the *Leptothorax tuborum* group (Hymenoptera: Formicidae). *Zeitschrift für Zoologische Systematik und Evolutionsforschung*, **29**, 165–175.
- Edwards AC, Zwarts L, Yamamoto A, Callaerts P, Mackay TFC (2009) Mutations in many genes affect aggressive behavior in *Drosophila melanogaster*. *BMC Biology*, **7**, 29.
- Evans JD, Wheeler DE (2000) Expression profiles during honeybee caste determination. *Genome Biology*, **2**, 1–6.
- Feldhaar H, Fiala B, Gadau J, Mohamed M, Maschwitz U (2003) Molecular phylogeny of *Crematogaster* subgenus *Decacrema* ants (Hymenoptera: Formicidae) and the colonization of *Macaranga* (Euphorbiaceae) trees. *Molecular Phylogenetics and Evolution*, **27**, 441–452.
- Feldhaar H, Foitzik S, Heinze J (2008) Lifelong commitment to the wrong partner: hybridization in ants. *Philosophical transactions of the Royal Society of London Series B: Biological Sciences*, **363**, 2891–2899.
- Gibson G, Weir B (2005) The quantitative genetics of transcription. *Trends in Genetics*, **21**, 616–623.
- Gräff J, Jemielity S, Parker JD, Parker KM, Keller L (2007) Differential gene expression between adult queens and workers in the ant *Lasius niger*. *Molecular Ecology*, **16**, 675–683.
- Graveley BR, Brooks AN, Carlson JW *et al.* (2011) The developmental transcriptome of *Drosophila melanogaster*. *Nature*, **471**, 473–479.
- Haerty W, Singh RS (2006) Gene regulation divergence is a major contributor to the evolution of Dobzhansky-Muller incompatibilities between species of *Drosophila*. *Molecular Biology and Evolution*, **23**, 1707–1714.
- Haldane JBS (1922) Sex ratio and unisexual sterility in hybrid animals. *Journal of Genetics*, **12**, 101–109.
- Helms Cahan S, Keller L (2003) Complex hybrid origin of genetic caste determination in harvester ants. *Nature*, **424**, 306–309.
- Helms Cahan S, Vinson SB (2003) Reproductive division of labor between hybrid and nonhybrid offspring in a fire ant hybrid zone. *Evolution*, **57**, 1562–1570.
- Helms Cahan S, Parker JD, Rissing SW *et al.* (2002) Extreme genetic differences between queens and workers in hybridizing Pogonomyrmex harvester ants. *Proceedings of the Royal Society B: Biological Sciences*, **269**, 1871–1877.
- Helms Cahan S, Julian GE, Rissing SW *et al.* (2004) Loss of phenotypic plasticity generates genotype-caste association in harvester ants. *Current Biology*, **14**, 2277–2282.
- Hey J, Kliman RM (1993) Population genetics and phylogenetics of DNA sequence variation at multiple loci within the *Drosophila melanogaster* species complex. *Molecular Biology and Evolution*, **10**, 804–822.
- Hoffman EA, Goodisman MAD (2007) Gene expression and the evolution of phenotypic diversity in social wasps. *BMC Biology*, **5**, 23.
- Hollocher H, Agopian K, Waterbury J, O'Neill RW, Davis AW (2000) Characterization of defects in adult germline development and oogenesis of sterile and rescued female hybrids in crosses between *Drosophila simulans* and *Drosophila melanogaster*. *The Journal of Experimental Zoology*, **288**, 205–218.
- Huang DW, Sherman BT, Lempicki RA (2009) Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols*, **4**, 44–57.
- Hunt BG, Goodisman MAD (2010) Evolutionary variation in gene expression is associated with dimorphism in eusocial vespid wasps. *Insect Molecular Biology*, **19**, 641–652.
- Hunt BG, Wyder S, Elango N *et al.* (2010) Sociality is linked to rates of protein evolution in a highly social insect. *Molecular Biology and Evolution*, **27**, 497–500.
- Hunt BG, Ometto L, Wurm Y *et al.* (2011) Relaxed selection is a precursor to the evolution of phenotypic plasticity. *Proceedings of the National Academy of Sciences of the United States of America*, **108**, 15936–15941.
- Jessen K, Klinkicht M (1990) Hybridization in the social parasitic ant genus *Epimyrma* (Hymenoptera, Formicidae). *Insectes Sociaux*, **37**, 273–293.
- Jouvenaz DP, Allen GE, Banks WA, Wojcik DP (1977) A survey for pathogens of fire ants, *Solenopsis* spp., in the Southeastern United States. *The Florida Entomologist*, **60**, 275–279.
- Judice CC, Carazzole MF, Festa F *et al.* (2006) Gene expression profiles underlying alternative caste phenotypes in a highly eusocial bee, *Melipona quadrifasciata*. *Insect Molecular Biology*, **15**, 33–44.
- Koevoets T, Niehuis O, van de Zande L, Beukeboom LW (2012) Hybrid incompatibilities in the parasitic wasp genus *Nasonia*: negative effects of hemizygoty and the identification of transmission ratio distortion loci. *Heredity*, **108**, 302–311.
- Kong EC, Allouche L, Chapot PA *et al.* (2010) Ethanol-regulated genes that contribute to ethanol sensitivity and rapid tolerance in *Drosophila*. *Alcoholism, Clinical and Experimental Research*, **34**, 302–316.
- Kulmuni J, Seifert B, Pamilo P (2010) Segregation distortion causes large-scale differences between male and female genomes in hybrid ants. *Proceedings of the National Academy of Sciences of the United States of America*, **107**, 7371–7376.
- Landry CR, Hartl DL, Ranz JM (2007) Genome clashes in hybrids: insights from gene expression. *Heredity*, **99**, 483–493.
- Linksvayer TA, Wade MJ (2009) Genes with social effects are expected to harbor more sequence variation within and between species. *Evolution*, **63**, 1685–1696.
- Mavarez J, Audet C, Bernatchez L (2009) Major disruption of gene expression in hybrids between young sympatric anadromous and resident populations of brook charr (*Salvelinus fontinalis* Mitchell). *Journal of Evolutionary Biology*, **22**, 1708–1720.
- Mcdermott SR, Kliman RM (2008) Estimation of isolation times of the island species in the *Drosophila simulans* complex from multilocus DNA sequence data. *PLoS ONE*, **3**, e2442.

- Meiklejohn CD, Townsend JP (2006) A Bayesian method for analysing spotted microarray data. *Briefings in Bioinformatics*, **6**, 318–330.
- Michalak P, Noor MAF (2003) Genome-wide patterns of expression in *Drosophila* pure species and hybrid males. *Molecular Biology and Evolution*, **20**, 1070–1076.
- Moehring AJ, Teeter KC, Noor MAF (2007) Genome-wide patterns of expression in *Drosophila* pure species and hybrid males. II. Examination of multiple-species hybridizations, platforms, and life cycle stages. *Molecular Biology and Evolution*, **24**, 137–145.
- Muller H (1942) Isolating mechanisms, evolution and temperature. *Biological Symposia*, **6**, 71–125.
- Nehrt NL, Clark WT, Radivojac P, Hahn MW (2011) Testing the ortholog conjecture with comparative functional genomic data from mammals. *PLoS Computational Biology*, **7**, e1002073.
- Nygaard S, Zhang G, Schiøtt M *et al.* (2011) The genome of the leaf-cutting ant *Acromyrmex echinator* suggests key adaptations to advanced social life and fungus farming. *Genome Research*, **21**, 1339–1348.
- Ometto L, Shoemaker D, Ross KG, Keller L (2011) Evolution of gene expression in fire ants: the effects of developmental stage, caste, and species. *Molecular Biology and Evolution*, **28**, 1381–1392.
- Orr H (1993a) A mathematical model of Haldane's rule. *Evolution*, **47**, 1606–1611.
- Orr HA (1993b) Haldane's rule has multiple genetic causes. *Nature*, **361**, 532–533.
- Ortiz-Barrientos D, Counterman BA, Noor MAF (2007) Gene expression divergence and the origin of hybrid dysfunctions. *Genetica*, **129**, 71–81.
- Pearson B (1983) Hybridisation between the ant species *Lasius niger* and *Lasius alienus*: the genetic evidence. *Insectes Sociaux*, **30**, 402–411.
- Pereboom JJM, Jordan WC, Sumner S, Hammond RL, Bourke AFG (2005) Differential gene expression in queen-worker caste determination in bumble-bees. *Proceedings of the Royal Society of London Series B: Biological Sciences*, **272**, 1145–1152.
- Pusch K, Heinze J, Foitzik S (2006) The influence of hybridization on colony structure in the ant species *Temnothorax nylander* and *T. crassispinus*. *Insectes Sociaux*, **53**, 439–445.
- R Development Core Team (2009) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Ranz JM, Namgyal K, Gibson G, Hartl DL (2004) Anomalies in the expression profile of interspecific hybrids of *Drosophila melanogaster* and *Drosophila simulans*. *Genome Research*, **14**, 373–379.
- Renaut S, Bernatchez L (2011) Transcriptome-wide signature of hybrid breakdown associated with intrinsic reproductive isolation in lake whitefish species pairs (*Coregonus* spp. Salmonidae). *Heredity*, **106**, 1003–1011.
- Renaut S, Nolte AW, Bernatchez L (2009) Gene expression divergence and hybrid misexpression between lake whitefish species pairs (*Coregonus* spp. Salmonidae). *Molecular Biology and Evolution*, **26**, 925–936.
- Riddle NC, Birchler JA (2003) Effects of reunited diverged regulatory hierarchies in allopolyploids and species hybrids. *Trends in Genetics*, **19**, 597–600.
- Rockman MV, Kruglyak L (2006) Genetics of global gene expression. *Nature Reviews Genetics*, **7**, 862–872.
- Ross KG, Robertson JL (1990) Developmental stability, heterozygosity, and fitness in two introduced fire ants (*Solenopsis invicta* and *S. richteri*) and their hybrid. *Heredity*, **64**, 93–103.
- Ross KG, Shoemaker DD (2005) Species delimitation in native South American fire ants. *Molecular Ecology*, **14**, 3419–3438.
- Ross KG, Trager JC (1990) Systematics and population genetics of fire ants (*Solenopsis saevissima* Complex) from Argentina. *Evolution*, **44**, 2113–2134.
- Ross KG, vander Meer RK, Fletcher DJC, Vargo EL (1987) Biochemical phenotypic and genetic studies of two introduced fire ants and their hybrid (Hymenoptera: Formicidae). *Evolution*, **41**, 280–293.
- Scharf ME, Wu-Scharf D, Pittendrigh BR, Bennett GW (2003) Caste- and development-associated gene expression in a lower termite. *Genome Biology*, **4**, R62.
- Schilthuizen M, Giesbers MCWG, Beukeboom LW (2011) Haldane's rule in the 21st century. *Heredity*, **107**, 95–102.
- Schluter D (2001) Ecology and the origin of species. *Trends in Ecology & Evolution*, **16**, 372–380.
- Schwander T, Helms Cahan S, Keller L (2006) Genetic caste determination in *Pogonomyrmex* harvester ants imposes costs during colony founding. *Journal of Evolutionary Biology*, **19**, 402–409.
- Seifert B (1999) Interspecific hybridisations in natural populations of ants by example of a regional fauna (Hymenoptera, Formicidae). *Insectes Sociaux*, **46**, 45–52.
- Sen Sarma M, Whitfield CW, Robinson GE (2007) Species differences in brain gene expression profiles associated with adult behavioral maturation in honey bees. *BMC Genomics*, **8**, 202.
- Shoemaker DD, Ross KG, Arnold ML (1996) Genetic structure and evolution of a fire ant hybrid zone. *Evolution*, **50**, 1958–1976.
- Shoemaker DD, Ahrens ME, Ross KG (2005) Molecular phylogeny of fire ants of the *Solenopsis saevissima* species-group based on mtDNA sequences. *Molecular Phylogenetics and Evolution*, **38**, 200–215.
- Sumner S, Pereboom JJM, Jordan WC (2006) Differential gene expression and phenotypic plasticity in behavioural castes of the primitively eusocial wasp, *Polistes canadensis*. *Proceedings of the Royal Society of London Series B: Biological Sciences*, **273**, 19–26.
- Taylor EB, Gerlinsky C, Farrell N, Gow JL (2012) A test of hybrid growth disadvantage in wild, free-ranging species pairs of threespine stickleback (*Gasterosteus aculeatus*) and its implications for ecological speciation. *Evolution*, **66**, 240–251.
- Trager J (1991) A revision of the fire ants, *Solenopsis geminata* group (Hymenoptera, Formicidae, Myrmicinae). *Journal of the New York Entomological Society*, **99**, 141–198.
- True JR, Haag ES (2001) Developmental system drift and flexibility in evolutionary trajectories. *Evolution & Development*, **3**, 109–119.
- Tschinkel W (2006) *The Fire Ants*. Harvard University Press, Cambridge, Mass.
- Turelli M, Orr H (1995) The dominance theory of Haldane's rule. *Genetics*, **140**, 389–402.

- Tweedie S, Ashburner M, Falls K *et al.* (2009) FlyBase: enhancing *Drosophila* Gene Ontology annotations. *Nucleic Acids Research*, **37**, D555–D559.
- Umphrey GJ (2006) Sperm parasitism in ants: selection for interspecific mating and hybridization. *Ecology*, **87**, 2148–2159.
- Wang J, Jemielity S, Uva P *et al.* (2007) An annotated cDNA library and microarray for large-scale gene-expression studies in the ant *Solenopsis invicta*. *Genome Biology*, **8**, R9.
- Weil T, Korb J, Rehli M (2009) Comparison of queen-specific gene expression in related lower termite species. *Molecular Biology and Evolution*, **26**, 1841–1850.
- Wijnen H, Naef F, Boothroyd C, Claridge-Chang A, Young MW (2006) Control of daily transcript oscillations in *Drosophila* by light and the circadian clock. *PLoS Genetics*, **2**, e39.
- Wurm Y, Uva P, Ricci F *et al.* (2009) Fourmidable: a database for ant genomics. *BMC Genomics*, **10**, 5.
- Wurm Y, Wang J, Riba-Grognuz O *et al.* (2011) The genome of the fire ant *Solenopsis invicta*. *Proceedings of the National Academy of Sciences of the United States of America*, **108**, 5679–5684.
- Wurmser F, Ogereau D, Mary-Huard T *et al.* (2011) Population transcriptomics: insights from *Drosophila simulans*, *Drosophila sechellia* and their hybrids. *Genetica*, **139**, 465–477.

L. Ometto's main research focus is the molecular basis of adaptation. This work is part of his post-doctoral project on the evolution of gene expression in fire ants. K. Ross's research focuses on the evolutionary genetics of social Hymenoptera and other insects, with much of his work concerning native and introduced fire ant populations. The main research interest of D. Shoemaker is the population and evolutionary genetics of various insects, focusing on fire ants and Wolbachia-insect interactions. L. Keller works on various aspects of evolutionary ecology and social behavior in ants.

Data accessibility

Microarray data are deposited in NCBI's Gene Expression Omnibus database under accession number GSE35217.

BAGEL output data is deposited in the Dryad repository: doi:10.5061/dryad.m0r5qv24.

Supporting information

Additional supporting information may be found in the online version of this article.

Fig. S1 Principal components analysis of gene expression levels. Three-letter labels identify taxonomic group (h = hybrids; i = *S. invicta*; r = *S. richteri*), caste (M = males; Q = queens; W = workers), and developmental stage (p = pupae; a = adults). Percentages of variability explained by each component are shown in parentheses.

Fig. S2a Venn diagrams depicting numbers of genes over-expressed in pupal males, workers, and queens of hybrids (hyb), *S. invicta* (inv), and *S. richteri* (ric). Overlapping areas indicate differentially expressed genes common to two or three of the taxonomic groups. Circle size is proportional to the number of genes.

Fig. S2b Venn diagrams depicting numbers of genes over-expressed in adult males, workers, and queens of hybrids (hyb), *S. invicta* (inv), and *S. richteri* (ric). See Fig. S2a (Supporting information) caption for additional explanation.

Fig. S3a Venn diagrams depicting numbers of genes under-expressed in pupal males, workers, and queens of hybrids (hyb), *S. invicta* (inv), and *S. richteri* (ric). See Fig. S2a (Supporting information) caption for additional explanation.

Fig. S3b Venn diagrams depicting numbers of genes under-expressed in adult males, workers, and queens of hybrids (hyb), *S. invicta* (inv), and *S. richteri* (ric). See Fig. S2a (Supporting information) caption for additional explanation.

Table S1. Complete list of the microarray hybridizations performed. Each hybridization contrasted the expression levels in the sample of interest with those in a common reference.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.